





Search PubMed	for Tsichlis P AND Gfi-1 Go Clear
rearen	gimits Preside that the Cippoard Details
	Display Summary Show: 20 Sort Send to Text
iextilers m	Items 1-7 of 7 One page.
Entrez Publika Durweu Haya Kalu Mew Notes on y Hayakhay	1:Zhu J. Guo F. Min B. Watson CJ. Hu-Li J. Young HA. Tsichlis PN. Related Actors of the Paul WE.  Growth factor independent-1 induced by IL-4 regulates Th2 cell proliferation.  Immunity, 2002 May:16(5):733-44.  PMID: 12049724 [PubMed - indexed for MLDLINE]
riebuided Selvices dournals Catomy ( Me Shiphous Single Catom Matimal Decomposition ( Matimal	2: Long B. Grimes HE., Yang TY, Beat SF, Qin Z, Du K, Fl-Deiry WS. Apparent Art new Lonks Islandis PN.  The Gfi-1B proto-oncoprotein represses p21WAF1 and inhibits myeloid cell differentiation.  Mol Cell Biol. 1998 May 18(5):216?-13  PMID: 9566867 [PubMed - indexed for MFDLINE]
Related Resources Order Documents NLM Gatoway TOXNET Consumer assam Changa Alerts	The Gfi-1 protooncoprotein represses Bax expression and inhibits T-cell death.  Proc Natl Acad Sci U.S.A. 1996 Dec 10,93(25):14569-73.  PMID: 8962093 [PubMed - indexed for MEDLINE]
Classian in a notice of the contract of the co	4:Grimes HL, Chan TO, Zweidler-McKay PA, Tong B, Tsichlis PN Related Articles Links
Prvacy runc.	The Gfi-1 proto-oncoprotein contains a novel transcriptional repressor domain. SNAG, and inhibits G1 arrest induced by interleukin-2 withdrawal. Mol Cell Biol. 1996 Nov:16(11):6263-72.  PMID: 8887656 [PubMed - indexed for MEDLINE]
	Gti-1 encodes a nuclear zinc finger protein that binds DNA and functions as a transcriptional repressor.  Mol Cell Biol 1996 Aug 16(8):4024-34  PMID: 8754800 [PubMed - indexed for MEDLINE]
	Prolactin (PRL)-dependent expression of a zinc finger protein-encoding gene. Gff-1, in Nb2 lymphoma cells: constitutive expression in autonomous sublines.  Indocrmology 1995 Apr. 136(4) 1805-8 PMID 7898694 [PubMed - indexed for MI DEINE]
	□ 7: 1. But a State of the control of the Progression of interleukin/2 (II -2)-dependent rat T cell lymphoma lines to

II.-2-independent growth following activation of a gene (Gfi-1) encoding a novel zinc finger protein.

Mol Cell Biol (1993 Mar. 13(3):1759-68.

PMID: 8441411 [PubMed - indexed for MEDLINE]

Display Summary Show 20 Sort Send to Text Items 1-7 of 7 One page.

Or te to the Help Desk
NOB\* | NLM | N\*H
Department of Health & Human Services
Freedom of Information Act | Disclaimer







Search PubMed	for Grimes HL AND Gfi-1 Go Clear
	and tell in Proposition of the Proposition of the Details
***	
	Display Summary → Show: 20 → Sort → Send to Text →
* x *	Items 1-5 of 5 One page.
Enter <sup>e</sup> l diller Succes	1: Long B. Grimes HL, Yang TY, Bear SL, Qin Z, Du K, El-Deiry WS. Related Articles Times Tsichlis PN.
k <del>in</del> g kabu	The Gfi-1B proto-oncoprotein represses p21WAF1 and inhibits myeloid cell
New York Con-	differentiation. Mol Cell Biol. 1998 May:18(5):2462-73. PMID: 9566867 [PubMed   indexed for MEDLINE]
위ut Med 3년 vires 과과마왕동 widthse	<b>2:</b> Grimes III., Gilks CB, Chan TO, Porter S, Tsiehlis PS. Related Actions Links
theSelf Energy in Matcher Single Citation Matcher Baton Citation Matcher	The Gfi-1 protooncoprotein represses Bax expression and inhibits T-cell death.
	Proc Natl Acad Sci U.S.A. 1996 Dec 10:93(25):14569-73. PMID: 8962093 [PubMed - indexed for MEDLINE]
Relatert for Social Soc	3:Grimes HL, Chan TO, Zweidler-McKay PA, Tong B, Tsichlis PN. Related Articles 1 - ks
	The Gfi-1 proto-oncoprotein contains a novel transcriptional repressor domain. SNAG, and inhibits G1 arrest induced by interleukin-2 withdrawal Mol Cell Biol. 1996 Nov:16(11):6263-72 PMID: 8887656 [PubMed - indexed for MEDLINE]
Olmgos Arents Olmgos Traxellador	
Purifice to the time	4: Zweidler-Mckay PA, Grimes HL, Flubacher MM, Tsichlis PN Related Articles Links
	Gfi-1 encodes a nuclear zinc finger protein that binds DNA and functions as a transcriptional repressor.  Mol Cell Biol. 1996 Aug;16(8):4024-34.  PMID: 8754800 [PubMed - indexed for MEDLINE]
	☐ 5: Onlies C.B. Beut Sir, Origines III : Ts.eldis P.S. Related Articles (Links
	Progression of interleukin-2 (IL-2)-dependent rat T cell lymphoma lines to IL-2-independent growth following activation of a gene (Gfi-1) encoding a novel zinc finger protein.  Mol Cell Biol. 1993 Mar;13(3):1759-68.  PMID: 8441411 [PubMed - indexed for MEDLINE]
	Display Summary - Show 20 - Sort - Send to Text -
	Items 1-5 of 5 One page.

Dife to the Help Nesk Tile 1, 10 1, 10 1, 1 Tile arthern if Health & Human Services Freedom of information Act Disclaimer







Search PubMed	Tor Zweidler-McKay P AND Gfi-1 Go Clear
	umits Preseduntado History Cippulata Bétals
At the end of	
	Display Summary → Show: 20 → Sort → Send to Text →
Text sers to	Items 1-2 of 2 One page
동시() 등 () 1 · 기기 ()	1:Orimes HL, Chan TO, Zweidler-McKay PA, Tong B, Tsichlis PN, Related Httldes Link
in to the	The Gfi-1 proto-oncoprotein contains a novel transcriptional repressor
· · ·	domain, SNAG, and inhibits G1 arrest induced by interleukin-2 withdrawal.
New Notes of V	Mol Cell Biol. 1996 Nov:16(11):6263-72. PMID: 8887656 (PubMed - indexed for MEDLINE)
Publica 80 vites aumum a	2://weidler-Mckay PA, Grimes HL, Flubacher MM, Tsichlis PN  Gffi-1 encodes a nuclear zinc finger protein that binds DNA and functions as a transcriptional repressor.  Mol Cell Biol. 1996 Aug;16(8):4024-34.  PMID: 8754800 [PubMed - indexed for MEDLINE]
Related Resources Index Decisions NUM Cottonsy TOXII Consumer means If the first the Public Learns	

Write to the Help Desk NCBI | NLM | NiH Department of Health & Human Services Freedom of Information Act | Disclaimer